

Asthma Action Plan for

Modified by Dr Mike Levin from an action plan used at the Royal Hospital for sick children, Edinburgh.



Tel: 0861 ASTHMA (278462)

Fax: 086 655 0809

naepr@netactive.co.za

www.asthma.co.za

- Take your controller medication every day whether you feel well or unwell.
- Visit the Doctor /Asthma Clinic twice a year even if asthma is well controlled.
- Take your medication / pumps / spacers with you to every doctors / nurses visit.
- Take this plan to each visit so it can be updated.
- Take the symptom or peak flow diary to each visit.

Doctors Phone No.

Hospital Phone No.

Date

Normal Peak Flow

Best Peak Flow

Asthma sufferers can:

- Have NO Symptoms.
- Have a normal lifestyle, play sport and sleep well.
- Have as few acute attacks as possible.
- Miss little or no school and work.
- Have your best possible peak flow.

What are the three zones?

Green Zone: *Your asthma is under control. This is where you want to be most of the time.*

Orange Zone - Caution: *Your asthma is not under control. The medication may need to be changed. Follow the advice in this plan and keep a symptom and medication diary. Make an appointment to see your doctor or asthma nurse.*

Red Zone - Red Alert: *Your asthma is critical! Follow the Red Zone Action and see a doctor immediately or go to the closest emergency room.*

GREEN ZONE - GO

Asthma is under control when:

- No cough or wheeze.
- Can play games and sport normally.
- No sleep disturbance.
- Using reliever less than 3 times a week.

AND

Peak flows are greater than 80%

ACTION: Take normal medicines

1. Controller

Strength

Your device is

Take puffs

When: every day

2. Other medicines

Medicine

Dose

When

.....

3. Reliever

Device

Take puffs as required

And if necessary take puffs

10 - 15 minutes before sports and activity.

ORANGE ZONE - CAUTION

Asthma is getting worse if there is:

- Cough, wheeze or tight chest.
- Waking at night with asthma symptoms.
- Need to use the reliever inhaler more than 3 times a week.
- Problems playing or doing sport.

OR

Peak flow recordings are between (50%) and (80%)

ACTION: Take normal medicines AND

Increase the reliever inhaler

to puffs four times a day until you are back in the Green zone.

Continue to take your controller inhaler as normal to prevent your symptoms

Other action:

.....

.....

.....

.....

If there is no improvement make an appointment to see your Doctor or Asthma Nurse. Fill in a symptom and medication diary every day and take it with you to the Dr or Asthma Nurse.

RED ZONE - ALERT

Asthma is dangerous when:

- Breathing is hard and fast
- Can't talk easily or feed easily
- Severe shortness of breath
- The reliever pump is not helping

OR

Peak flow is below 50%

ACTION: Call an ambulance or go to a doctor NOW, even if symptoms get better!

Take 1 puff of reliever every minute for 10 minutes. Use a spacer if you have one.

Repeat this if there is no improvement as often as you need.

While waiting: Give 1 puff of reliever every minute for 10 minutes using a spacer if you have one.

- Use steroid tablets or syrup.
Your dose is
- Keep calm
- Sit up to help breathe
- Loosen clothing